

1. The human eye can focus on objects at different distances by adjusting the focal length of the eye lens. This is due to



**Ans.** (b) The human eye can change the focal length of the eye lens to see the objects situated at various distances from the eye. This is possible due to the power of accommodation of the eye lens.



**Ans.** (d) The human eye forms the image of an object at its retina.



**Ans.** (c) The least distance of distinct vision is an object's minimum distance to see a clear and distinct image. It is 25 cm for a young adult with normal vision.



**Ans.** (c) The relaxation or contraction of ciliary muscles changes the curvature of the eye lens. The change in curvature of the eye lens changes the focal length of the eyes. Hence, the change in focal length of an eye lens is caused by the action of ciliary muscles.

- 5.** A person needs a lens of power  $-5.5$  dioptres for correcting his distant vision. For correcting his near vision, he needs a lens of power  $+1.5$  dioptre. What is the focal length of the lens required for correcting (i) distant vision and (ii) near vision?

**Ans.** For distant vision  $\equiv -0.181$  m, for near vision  $\equiv 0.667$  m

The power  $P$  of a lens of focal length  $f$  is given by the relation

$$P = \frac{1}{f(\text{in meters})}$$

(i) Power of the lens used for correcting distant vision =  $-5.5$  D

$$\text{The focal length of the required lens, } f = \frac{1}{p}$$

$$f = \frac{1}{-5.5} = -0.181 \text{ m}$$

The focal length of the lens for correcting distant vision is  $-0.181$  m.

(ii) Power of the lens used for correcting near vision =  $+1.5$  D

$$\text{The focal length of the required lens, } f = \frac{1}{p}$$

$$f = \frac{1}{1.5} = +0.667 \text{ m}$$

The focal length of the lens for correcting near vision is  $0.667$  m.

**6.** The far point of a myopic person is  $80$  cm in front of the eye. What is the nature and power of the lens required to correct the problem?

**Ans.** The person is suffering from an eye defect called myopia. In this defect, the image is formed in front of the retina. Hence, a concave lens is used to correct this defect of vision.

Object distance,  $u = \text{infinity} = \infty$

Image distance,  $v = -80$  cm

Focal length =  $f$

According to the lens formula,

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

$$-\frac{1}{80} - \frac{1}{\infty} = \frac{1}{f}$$

$$\frac{1}{f} = -\frac{1}{80}$$

$$f = -80 \text{ cm} = -0.8 \text{ m}$$

We know,

$$\text{Power, } P = \frac{1}{f \text{ (in meters)}}$$

$$P = \frac{1}{-0.8} = -1.25$$

The person requires a concave lens of power  $-1.25$  D to correct his defect.

**7.** Why is a normal eye not able to see the objects placed closer than  $25$  cm?

**Ans.** A normal eye cannot see the objects placed closer than  $25$  cm because the eyes' ciliary muscles are unable to contract beyond a certain limit.


If the object is placed at a distance less than 25 cm from the eye, then the object appears blurred and produces strain in the eyes.

**8.** What happens to the image distance in the eye when we increase the distance of an object from the eye?

**Ans.** Since the size of the eyes cannot increase or decrease, the image distance remains constant. When we increase the distance of an object from the eye, the image distance in the eye does not change. The increase in the object distance is compensated by the change in the focal length of the eye lens. The focal length of the eyes changes so that the image is always formed at the retina of the eye.

**9.** Make a diagram to show how hypermetropia is corrected. The near point of a hypermetropic eye is 1 m. What is the power of the lens required to correct this defect? Assume that the near point of the normal eye is 25 cm.

**Ans.** A person suffering from hypermetropia can see distinct objects clearly but faces difficulty in seeing nearby objects. It happens because the eye lens focuses on the incoming divergent rays beyond the retina. This defect of vision is corrected by using a convex lens. A convex lens of suitable power converges the incoming light so that the image is formed on the retina, as shown in the following figure.



Correction for hypermetropic eye

The convex lens creates a virtual image of a nearby object ( $N'$  in the figure) at the near point of vision ( $N$ ) of the person suffering from hypermetropia.

The given person will be able to see the object kept at 25 cm (near the point of the normal eye) if the image of the object is formed at his near point, which is given as 1 m.

Object distance,  $u = -25 \text{ cm}$

Image distance,  $v = -1 \text{ m} = -100 \text{ cm}$

Focal length,  $f$

Using the lens formula,

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f} \quad \frac{1}{-100} - \frac{1}{-25} = \frac{1}{f}$$

$$\frac{1}{f} = \frac{1}{25} - \frac{1}{100}$$

$$\frac{1}{f} = \frac{4-1}{100}$$

$$f = \frac{100}{3} = 33.3 \text{ cm} = 0.33 \text{ m}$$

$$\text{Power, } P = \frac{1}{f(\text{in meters})} = \frac{1}{0.33 \text{ m}} = + 3.0 \text{ D}$$

A convex lens of power +3.0 D is required to correct the defect.

**10.** Why does the sky appear dark instead of blue to an astronaut?

**Ans.** The sky appears dark instead of blue to an astronaut because no atmosphere can scatter the sunlight in outer space. As the sunlight is not scattered, no scattered light reaches the astronauts' eyes, and the sky appears black to them.

**11.** Why do stars twinkle?

**Ans.** Stars emit their light, and they twinkle due to the atmospheric refraction of light. Stars are very far away from the earth. Hence, they are considered as point sources of light. When the light coming from stars enters the earth's atmosphere, it gets refracted at different levels because of the variation in the air density at different levels of the atmosphere. When the starlight refracted by the atmosphere comes more towards us, it appears brighter than when it comes less towards us. Therefore, it appears as if the stars are twinkling at night.

**12.** Explain why the planets do not twinkle?

**Ans.** Planets do not twinkle because they appear larger than the stars as they are relatively closer to earth. Planets can be considered as a collection of a large number of point-size sources of light. The different parts of these planets produce either brighter or dimmer effects. The average of brighter and dimmer effects is zero. Hence, the twinkling effects of the planets are nullified, and they do not twinkle.

**13.** Why does the Sun appear reddish early in the morning?

**Ans.** During sunrise, the light rays coming from the Sun have to travel a greater distance in the earth's atmosphere before reaching our eyes. In this journey, the shorter wavelengths of lights are scattered out, and only longer wavelengths can reach our eyes. Since blue colour has a shorter wavelength and red colour has a longer wavelength, the red colour can reach our eyes after the atmospheric scattering of light. Therefore, the Sun appears reddish early in the morning.