




SpeedLabs

MATHS

CBSE 9th

TEEVRA EDUTECH PVT. LTD.

Q.1 In figure, ABCD is a parallelogram. $AE \perp DC$ and $CF \perp AD$. If $AB = 16$ cm, $AE = 8$ cm and $CF = 10$ cm, find AD.



Ans. ABCD is a parallelogram.

$$\therefore DC = AB \Rightarrow DC = 16 \text{ cm}$$

$AE \perp DC$ [Given]

Now Area of parallelogram ABCD = Base x Corresponding height

$$= DC \times AE = 16 \times 8 = 128^2$$

Using base AD and height CF, we can find,

Area of parallelogram


$$\Rightarrow 128 = AD \times 10$$

$$AD = \frac{128}{10} = 12.8 \text{ cm}$$

Q.2 If E, F, G and H are respectively the mid-points of the sides of a parallelogram ABCD,

show that $\text{ar}(\text{EFGH}) = \frac{1}{2} \text{ar}(\text{ABCD})$.

Ans. Given: A parallelogram ABCD. E, F, G and H are mid-points of AB, BC, CD and DA respectively.



To prove: $\text{ar}(\text{EFGH}) = \frac{1}{2} \text{ar}(\text{ABCD})$

Construction: Join HF

Proof: $\text{ar}(\Delta \text{GHF}) = \frac{1}{2} \text{ar}(\text{ll gm HFCD}) \dots\dots\dots \text{(i)}$

And $\text{ar}(\Delta \text{HEF}) = \frac{1}{2} \text{ar}(\text{ll gm HABF}) \dots\dots\dots \text{(ii)}$

[If a triangle and a parallelogram are on the same base and between the same parallel then the area of triangle is half of area of parallelogram]

Adding eq. (i) and (ii),


$$\text{ar}(\Delta GHF) + \text{ar}(\Delta HEF)$$

$$\frac{1}{2} = \text{ar}(\parallel \text{ gm HFCD}) + \frac{1}{2} \text{ar}(\parallel \text{ gm HABF})$$

$$\text{ar}(\parallel \text{ gm HEFG}) = \frac{1}{2} \text{ar}(\parallel \text{ gm ABCD})$$

Q.3 P and Q are any two points lying on the sides DC and AD respectively of a parallelogram ABCD. Show that $\text{ar}(\Delta APB) = \text{ar}(\Delta BQC)$.

Ans. Given: ABCD is a parallelogram. P is a point on DC and Q is a point on AD.



To prove: $\text{ar}(\Delta APB) = \text{ar}(\Delta BQC)$

Construction: Draw $PM \parallel BC$ and $QN \parallel DC$.

Proof: Since QC is the diagonal of parallelogram QNCD.

$$\therefore \text{ar}(\Delta QNC) = \frac{1}{2} \text{ar}(\parallel \text{ gm QNCD}) \dots\dots\dots(i)$$

Again, BQ is the diagonal of parallelogram ABNQ.

$$\therefore \text{ar}(\Delta BQN) = \frac{1}{2} \text{ar}(\parallel \text{ gm ABNQ}) \dots\dots\dots(ii)$$

Adding eq. (i) and (ii),

$$\begin{aligned} &\text{ar}(\Delta QNC) + \text{ar}(\Delta BQN) \\ &= \frac{1}{2} \text{ar}(\parallel \text{ gm QNCD}) + \frac{1}{2} \text{ar}(\parallel \text{ gm ABNQ}) \end{aligned}$$

$$\Rightarrow \text{ar}(\Delta BQC) = \frac{1}{2} \text{ar}(\parallel \text{ gm ABCD}) \dots\dots\dots(iii)$$

Again, AP is the diagonal of gm AMPD.

$$\therefore \text{ar}(\Delta APM) = \text{ar}(\parallel \text{ gm AMPD}) \dots\dots\dots(iv)$$

And PB is the diagonal of $\parallel \text{ gm PCBM}$.

$$\therefore \text{ar}(\Delta PBM) = \frac{1}{2} \text{ar}(\parallel \text{ gm PCBM}) \dots\dots\dots(v)$$

Adding eq. (iv) and (v),

$$\text{ar}(\Delta APM) + \text{ar}(\Delta PBM)$$


$$\frac{1}{2} = \text{ar}(\parallel \text{gm AMPD}) + \frac{1}{2} \text{ar}(\parallel \text{gm PCBM})$$

$$\Rightarrow \text{ar}(\Delta APB) = \frac{1}{2} \text{ar}(\parallel \text{gm ABCD}) \dots\dots\dots(\text{vi})$$

From eq. (iii) and (vi),

$$\text{ar}(\Delta BQC) = \text{ar}(\Delta APB) \text{ or } \text{ar}(\Delta APB) = \text{ar}(\Delta BQC)$$


Q.4 In figure, P is a point in the interior of a parallelogram ABCD Show that:



(i) $\text{ar}(\Delta APB) + \text{ar}(\Delta PCD) = \frac{1}{2} \text{ar}(\text{ABCD})$

(ii) $\text{ar}(\Delta APD) + \text{ar}(\Delta PBC) = \text{ar}(\Delta APB) + \text{ar}(\Delta PCD)$

Ans. (i) Draw a line passing through point P and parallel to AB which intersects AD at Q and BC at R respectively. Now ΔAPB and parallelogram ABRQ are on the same base AB and between same parallels AB and QR.



$$\therefore \text{ar}(\Delta APB) = \frac{1}{2} \text{ar}(\parallel \text{gm ABRQ}) \dots\dots\dots (\text{i})$$

Also ΔPCD and parallelogram DCRQ are on the same base AB and between same parallels AB and QR.

$$\therefore \text{ar}(\Delta PCD) = \frac{1}{2} \text{ar}(\parallel \text{gm DCRQ}) \dots\dots\dots (\text{ii})$$

Adding eq. (i) and (ii),

$$\text{ar}(\Delta APB) + \text{ar}(\Delta PCD)$$

$$= \frac{1}{2} \text{ar}(\parallel \text{gm ABRQ}) + \frac{1}{2} \text{ar}(\parallel \text{gm DCRQ})$$

$$\Rightarrow \text{ar}(\Delta APB) = \frac{1}{2} \text{ar}(\parallel \text{gm ABCD}) \dots\dots\dots (\text{iii})$$

(ii) Draw a line through P and parallel to AD which intersects AB at M and DC at N. Now ΔAPD and parallelogram AMND are on the same base AD and between same parallels AD and MN.

$$\therefore \text{ar} (\Delta APD) = \frac{1}{2} \text{ar} (\text{||gm AMND}) \dots\dots\dots (\text{iv})$$

Also, ΔPBC and parallelogram MNCB are on the same base BC and between same parallels BC and MN.

$$\therefore \text{ar} (\Delta PBC) = \frac{1}{2} \text{ar} (\text{|| gm MNCB}) \dots\dots\dots (\text{v})$$

Adding eq. (i) and (ii),

$$\begin{aligned} &\text{ar} (\Delta APD) + \text{ar} (\Delta PBC) \\ &= \frac{1}{2} \text{ar} (\text{|| gm AMND}) + \frac{1}{2} \text{ar} (\text{|| gm MNCB}) \end{aligned}$$

$$\Rightarrow \text{ar} (\Delta APD) = \frac{1}{2} \text{ar} (\text{|| gm ABCD}) \dots\dots\dots (\text{vi})$$


From eq. (iii) and (vi), we get,

$$\text{ar} (\Delta APB) + \text{ar} (\Delta PCD) = \text{ar} (\Delta APD) + \text{ar} (\Delta PBC)$$

$$\text{or } \text{ar} (\Delta APD) + \text{ar} (\Delta PBC) = \text{ar} (\Delta APB) + \text{ar} (\Delta PCD)$$

Hence proved.

Q.5 In figure, PQRS and ABRS are parallelograms and X is any point on the side BR. Show that:



(i) $\text{ar} (\text{PQRS}) = \text{ar} (\text{ABRS})$

(ii) $\text{ar} (\Delta \text{AXS}) = \frac{1}{2} \text{ar} (\text{PQRS})$

Ans. (i) Parallelogram PQRS and ABRS are on the same base SR and between the same parallels SR and PB.

$$\therefore \text{ar} (\text{|| gm PQRS}) = \frac{1}{2} \text{ar} (\text{|| gm ABRS}) \dots\dots\dots (\text{i})$$

[\therefore parallelograms on the same base and between the same parallels are equal in area]

(ii) ΔAXS and || gm ABRS are on the same base AS and between the same parallels AS and BR.


$$\therefore \text{ar} (\Delta \text{AXS}) = \frac{1}{2} \text{ar} (\text{|| gm ABRS}) \dots\dots\dots (\text{ii})$$

Using eq. (i) and (ii),

$$\text{ar} (\Delta AXS) = \frac{1}{2} \text{ar} (\parallel \text{gm PQRS})$$

Q.6 a farmer was having a field in the form of a parallelogram PQRS. She took any point A on RS and joined it to points P and Q. In how many parts the field is divided? What are the shapes of these parts? The farmer wants to sow wheat and pulses in equal portions of the field separately. How should she do it?

Ans. When A is joined with P and Q; the field is divided into three parts viz. ΔPAS , ΔAPQ and ΔAQR . ΔAPQ and parallelogram PQRS are on the same base PQ and between same parallels PQ and SR.



$$\therefore \text{ar} (\Delta APQ) = \frac{1}{2} \text{ar} (\parallel \text{gm PQRS})$$

It implies that triangular region APQ covers half portion of parallelogram shaped field PQRS.

So, if farmer sows wheat in triangular shaped field APQ then she will definitely sow pulses in other two triangular parts PAS and AQR.

Or

When she sows pulses in triangular shaped field APQ then she will sow wheat in other two triangular parts PAS and AQR.

