




SpeedLabs
Science

CBSE 10th

TEEVRA EDUTECH PVT. LTD.



Structure of a neuron

Functions of the three parts of a neuron.

Axon. It conducts messages away from the cell body.

Dendrite. It receives information from axon of another cell and conducts the messages towards the cell body.

Cell body. It contains nucleus, mitochondria, and other organelles. It is mainly concerned with the maintenance and growth.

6. How does phototropism occur in plants?

Ans. The growth movement in plants in response to light stimulus is known as phototropism. The shoots show positive phototropism and the roots show negative phototropism. This means that the shoots bend towards the source of light whereas the roots bend away from the light source.

Some examples of phototropism are as follows.

(a) The flower head of sunflower is positively phototropic and hence it moves from east to west along with the sun.

(b) The ovary stalk of groundnut is positively phototropic before fertilization and becomes negatively phototropic after fertilization, so that the fruit is formed underground.

7. Which signals will get disrupted in case of a spinal cord injury?

Ans. The reflex arc connections between the input and output nerves meet in a bundle in the spinal cord. In fact, nerves from all over the body meet in a bundle in the spinal cord on their way to the brain. In case of any injury to the spinal cord, the signals coming from the nerves as well as the signals coming to the receptors will be disrupted.

8. How does chemical coordination occur in plants?

Ans. In animals, control and coordination occur with the help of nervous system. However, plants do not have a nervous system.

Plants respond to stimuli by showing movements. The growth, development, and responses to the environment in plants is controlled and coordinated by a special class of chemical substances known as

hormones. These hormones are produced in one part of the plant body and are translocated to other needy parts. For example, a hormone produced in roots is translocated to other parts when required. The five major types of phytohormone are auxins, gibberellins, cytokinins, abscisic acid, and ethylene. These phytohormones are either growth promoters (such as auxins, gibberellins, cytokinins, and ethylene) or growth inhibitors such as abscisic acid.

9. What is the need for a system of control and coordination in an organism?

Ans. The maintenance of the body functions in response to changes in the body by working together of various integrated body systems is known as coordination. All the movements that occur in response to stimuli are carefully coordinated and controlled. In animals, the control and coordination movements are provided by nervous and muscular systems. The nervous system sends messages to and away from the brain. The spinal cord plays an important role in the relay of messages. In the absence of this system of control and coordination, our body will not be able to function properly. For example, when we accidentally touch a hot utensil, we immediately withdraw our hand. In the absence of nerve transmission, we will not withdraw our hand and may get burnt.

10. How are involuntary actions and reflex actions different from each other?

Ans. Involuntary actions cannot be consciously controlled. For example, we cannot consciously control the movement of food in the alimentary canal. These actions are however directly under the control of the brain. On the other hand, the reflex actions such as closing of eyes immediately when bright light is focused show sudden response and do not involve any thinking. This means that unlike involuntary actions, the reflex actions are not under the control of brain.

11. Compare and contrast nervous and hormonal mechanisms for control and coordination in animals.

Ans.

| | Nervous system mechanism | | Hormonal system mechanism |
|---|--|---|---|
| 1 | The information is conveyed in the form of electric impulse. | 1 | The information is conveyed in the form of chemical messengers. |
| 2 | The axons and dendrites transmit the information through a coordinated effort. | 2 | The information is transmitted or transported through blood. |
| 3 | The flow of information is rapid and the response is quick | 3 | The information travels slowly and the response is slow. |
| 4 | Its effects are short lived. | 4 | It has prolonged effects. |

12. What is the difference between the manner in which movement takes place in a sensitive plant and the movement in our legs?

Ans.

| | Movement in sensitive plants | | Movement in our legs |
|---|---|---|--|
| 1 | The movement that takes place in a sensitive plant such as Mimosa pudica occurs in response to touch (stimulus). | 1 | Movement in our legs is an example of voluntary actions |
| 2 | For this movement, the information is transmitted from cell to cell by electrochemical signals as plants do not have any specialized tissue for conduction of impulses. | 2 | The signal or messages for these actions are passed to the brain and hence are consciously controlled. |
| 3 | For this movement to occur, the plant cells change shape by changing the amount of water in them. | 3 | In animal muscle cells, some proteins are found which allow the movement to occur, |