



**CBSE 10<sup>th</sup>**

**TEEVRA EDUTECH PVT. LTD.**

# Quadratic Equation

## Exercise-4.2

**Q.1** Find the roots of the following quadratic equations by factorisation:  $x^2 - 3x - 10 = 0$

**Sol:**  $x^2 - 3x - 10$

$$\Rightarrow x^2 - 5x + 2x - 10$$

$$\Rightarrow x(x - 5) + 2(x - 5)$$

$$\Rightarrow (x - 5)(x + 2)$$

Roots of this equation are the values for which  $(x - 5)(x + 2) = 0$

$$\therefore x - 5 = 0 \text{ or } x + 2 = 0$$

i.e.,  $x = 5$  or  $x = -2$

**Q.2** Find the roots of the following quadratic equations by factorisation:  $2x^2 + x - 6 = 0$

**Sol:**  $2x^2 + x - 6$

$$\Rightarrow 2x^2 + 4x - 3x - 6$$

$$\Rightarrow 2x(x + 2) - 3(x + 2)$$

$$\Rightarrow (x + 2)(2x - 3)$$

Roots of this equation are the values for which  $(x + 2)(2x - 3) = 0$

$$\therefore x + 2 = 0 \text{ or } 2x - 3 = 0$$

i.e.,  $x = -2$  or  $x = 3/2$

**Q.3** Find the roots of the following quadratic equations by factorization:  $\sqrt{2x^2} + 7x + 5\sqrt{2}$

**Sol:**

$$\sqrt{2x^2} + 7x + 5\sqrt{2} = \sqrt{2x^2} + 5x + 2x + 5\sqrt{2} = x(\sqrt{2x} + 5) + \sqrt{2}(\sqrt{2x} + 5) = (\sqrt{2x} + 5)(x + \sqrt{2})$$

Root of this equation are the value for which  $(\sqrt{2x} + 5)(x + \sqrt{2}) = 0$

$$\therefore \sqrt{2x} + 5 = 0 \text{ or } x + \sqrt{2} = 0 \text{ i.e., } x = \frac{-5}{\sqrt{2}} \text{ or } x = -\sqrt{2}$$

**Q.4** Find the roots of the following quadratic equations by factorization:  $2x^2 - x + \frac{1}{8} = 0$

**Sol:**

$$= \frac{1}{8}(16x^2 - 8x + 1)$$

$$= \frac{1}{8}(16x^2 - 4x - 4x + 1)$$

$$= \frac{1}{8}(4x(4x - 1) - 1(4x + 1))$$

$$= \frac{1}{8}(4x - 1)^2$$

Roots of this equation are the values for which  $(4x - 1)^2 = 0$

Therefore,  $(4x - 1) = 0$  or  $(4x - 1) = 0$

$$\text{i.e., } x = \frac{1}{4} \text{ or } x = \frac{1}{4}$$

**Q.5** Find the roots of the following quadratic equations by factorisation:  $100x^2 - 20x + 1 = 0$

**Sol:**  $100x^2 - 20x + 1$

$$\Rightarrow 100x^2 - 10x - 10x + 1$$

$$\Rightarrow 10x(10x - 1) - 1(10x - 1)$$

$$\Rightarrow (10x - 1)^2$$

Roots of this equation are the values for which  $(10x - 1)^2 = 0$

Therefore,  $(10x - 1) = 0$  or  $(10x - 1) = 0$

$$\text{i.e., } x = \frac{1}{10} \text{ or } x = \frac{1}{10}$$

**Q.6** John and Jivanti together have 45 marbles. Both of them lost 5 marbles each, and the product of the number of marbles they now have is 124. Find out how many marbles they had to start with.

**Sol:** Let the number of John's marbles be  $x$ .

Therefore, number of Jivanti's marble =  $45 - x$

After losing 5 marbles,

Number of John's marbles =  $x - 5$

Number of Jivanti's marbles =  $45 - x - 5 = 40 - x$

It is given that the product of their marbles is 124.

$$\therefore (x - 5)(40 - x) = 124$$

$$\Rightarrow x^2 - 45x + 324 = 0$$

$$\Rightarrow x^2 - 36x - 9x + 324 = 0$$

$$\Rightarrow x(x - 36) - 9(x - 36) = 0$$

$$\Rightarrow (x - 36)(x - 9) = 0$$

Either  $x - 36 = 0$  or  $x - 9 = 0$

$$\text{i.e., } x = 36 \text{ or } x = 9$$

If the number of John's marbles = 36,

Then, number of Jivanti's marbles =  $45 - 36 = 9$

If number of John's marbles = 9,

Then, number of Jivanti's marbles =  $45 - 9 = 36$

**Q.7** A cottage industry produces a certain number of toys in a day. The cost of production of each toy(in rupees) was found to be 55 minus the number of toys produced in a day. On a particular day, the total cost of production was Rs 750. Find out the number of toys produced on that day.

**Sol:** Let the number of toys produced be  $x$ .

$\therefore$  Cost of production of each toy = Rs  $(55 - x)$

It is given that, total production of the toys = Rs 750

$$\therefore x(55 - x) = 750$$

$$\Rightarrow x^2 - 55x + 750 = 0$$

$$\Rightarrow x^2 - 25x - 30x + 750 = 0$$

$$\Rightarrow x(x - 25) - 30(x - 25) = 0$$

$$\Rightarrow (x - 25)(x - 30) = 0$$

Either  $x - 25 = 0$  or  $x - 30 = 0$

i.e.,  $x = 25$  or  $x = 30$

Hence, the number of toys will be either 25 or 30.

**Q.8** The following series is APs? If its a form an A.P. find the common difference  $d$  and write three more terms. 2, 4, 8, 16 ...

**Sol:** 2, 4, 8, 16 ...

It can be observed that

$$a_2 - a_1 = 4 - 2 = 2$$

$$a_3 - a_2 = 8 - 4 = 4$$

$$a_4 - a_3 = 16 - 8 = 8$$

i.e.,  $a_{k+1} - a_k$  is not the same every time. Therefore, the given numbers are not forming an A.P.

**Q.9** The following series is APs? If its a form an A.P. find the common difference  $d$  and write three more terms.  $2, \frac{5}{2}, 3, \frac{7}{2}, \dots$

**Sol:**

$$2, \frac{5}{2}, 3, \frac{7}{2}, \dots$$

It can be observed that

$$a_2 - a_1 = \frac{5}{2} - 2 = \frac{1}{2}$$

$$a_3 - a_2 = 3 - \frac{5}{2} = \frac{1}{2}$$

$$a_4 - a_3 = \frac{7}{2} - 3 = \frac{1}{2}$$

i.e.,  $a_{k+1} - a_k$  is same every time.

Therefore,  $d = \frac{1}{2}$  and the given numbers are in A.P.

Three more terms are

$$a_5 = \frac{7}{2} + \frac{1}{2} = 4$$

$$a_6 = 4 + \frac{1}{2} = \frac{9}{2}$$

$$a_7 = \frac{9}{2} + \frac{1}{2} = 5$$

**Q.10** The following series is APs? If its a form an A.P. find the common difference  $d$  and write three more terms.  $-1.2, -3.2, -5.2, -7.2 \dots$

**Sol:**  $-1.2, -3.2, -5.2, -7.2 \dots$

It can be observed that

$$a_2 - a_1 = (-3.2) - (-1.2) = -2$$

$$a_3 - a_2 = (-5.2) - (-3.2) = -2$$

$$a_4 - a_3 = (-7.2) - (-5.2) = -2$$

i.e.,  $a_{k+1} - a_k$  is same every time. Therefore,  $d = -2$

The given numbers are in A.P.

Three more terms are

$$a_5 = -7.2 - 2 = -9.2$$

$$a_6 = -9.2 - 2 = -11.2$$

$$a_7 = -11.2 - 2 = -13.2$$

**Q.11** The following series is APs? If its a form an A.P. find the common difference  $d$  and write three more terms.  $-10, -6, -2, 2 \dots$

**Sol:**  $-10, -6, -2, 2 \dots$

It can be observed that

$$a_2 - a_1 = (-6) - (-10) = 4$$

$$a_3 - a_2 = (-2) - (-6) = 4$$

$$a_4 - a_3 = (2) - (-2) = 4$$

i.e.,  $a_{k+1} - a_k$  is same every time. Therefore,  $d = 4$

The given numbers are in A.P.

Three more terms are

$$a_5 = 2 + 4 = 6$$

$$a_6 = 6 + 4 = 10$$

$$a_7 = 10 + 4 = 14$$

**Q.12** The following series is APs? If its a form an A.P. find the common difference d and write three more terms.  $3, 3 + \sqrt{2}, 3 + 2\sqrt{2}, 3 + 3\sqrt{2} \dots$

**Sol:**

$$3, 3 + \sqrt{2}, 3 + 2\sqrt{2}, 3 + 3\sqrt{2} \dots$$

It can be observed that

$$a_2 - a_1 = 3 + \sqrt{2} - 3 = \sqrt{2}$$

$$a_3 - a_2 = 3 + 2\sqrt{2} - 3 - \sqrt{2} = \sqrt{2}$$

$$a_4 - a_3 = 3 + 3\sqrt{2} - 3 - 2\sqrt{2} = \sqrt{2}$$

i.e.,  $a_{k+1} - a_k$  is same every time.

Therefore,  $d = \sqrt{2}$  and the given numbers are in A.P.

Three more terms are

$$a_5 = 3 + 3\sqrt{2} + \sqrt{2} = 3 + 4\sqrt{2}$$

$$a_6 = 3 + 4\sqrt{2} + \sqrt{2} = 3 + 5\sqrt{2}$$

$$a_7 = 3 + 5\sqrt{2} + \sqrt{2} = 3 + 6\sqrt{2}$$

**Q.13** The following series is APs? If its a form an A.P. find the common difference d and write three more terms.  $0.2, 0.22, 0.222, 0.2222 \dots$

**Sol:**  $0.2, 0.22, 0.222, 0.2222 \dots$

It can be observed that

$$a_2 - a_1 = 0.22 - 0.2 = 0.02$$

$$a_3 - a_2 = 0.222 - 0.22 = 0.002$$

$$a_4 - a_3 = 0.2222 - 0.222 = 0.0002$$

i.e.,  $a_{k+1} - a_k$  is not the same every time.



Therefore, the given numbers are not in A.P.

**Q.14** The following series is APs? If its a form an A.P. find the common difference  $d$  and write three more terms.  $0, -4, -8, -12 \dots$

**Sol:**  $0, -4, -8, -12 \dots$

It can be observed that

$$a_2 - a_1 = (-4) - 0 = -4$$

$$a_3 - a_2 = (-8) - (-4) = -4$$

$$a_4 - a_3 = (-12) - (-8) = -4$$

i.e.,  $a_{k+1} - a_k$  is same every time. Therefore,  $d = -4$

The given numbers are in A.P.

Three more terms are

$$a_5 = -12 - 4 = -16$$

$$a_6 = -16 - 4 = -20$$

$$a_7 = -20 - 4 = -24$$

**Q.15** The following series is APs? If its a form an A.P. find the common difference  $d$  and write three more terms.  $1, 3, 9, 27 \dots$

**Sol:**  $1, 3, 9, 27 \dots$

It can be observed that

$$a_2 - a_1 = 3 - 1 = 2$$

$$a_3 - a_2 = 9 - 3 = 6$$

$$a_4 - a_3 = 27 - 9 = 18$$

i.e.,  $a_{k+1} - a_k$  is not the same every time. Therefore, the given numbers are not in A.P.

**Q.16** The following series is APs? If its a form an A.P. find the common difference  $d$  and write three more terms.  $-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2} \dots$

**Sol:**

$$-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2} \dots$$

It can be observed that

$$a_2 - a_1 = \left(-\frac{1}{2}\right) - \left(-\frac{1}{2}\right) = 0$$

$$a_3 - a_2 = \left(-\frac{1}{2}\right) - \left(-\frac{1}{2}\right) = 0$$

$$a_4 - a_3 = \left(-\frac{1}{2}\right) - \left(-\frac{1}{2}\right) = 0$$

i.e.,  $a_{k+1} - a_k$  is same every time. Therefore,  $d = 0$

The given numbers are in A.P.

Three more terms are

$$a_5 = -\frac{1}{2} - 0 = -\frac{1}{2}$$

$$a_6 = -\frac{1}{2} - 0 = -\frac{1}{2}$$

$$a_7 = -\frac{1}{2} - 0 = -\frac{1}{2}$$

**Q.17** The following series is APs? If its a form an A.P. find the common difference  $d$  and write three more terms.  $a, 2a, 3a, 4a \dots$

**Sol:**  $a, 2a, 3a, 4a \dots$

It can be observed that

$$a_2 - a_1 = 2a - a = a$$

$$a_3 - a_2 = 3a - 2a = a$$

$$a_4 - a_3 = 4a - 3a = a$$

i.e.,  $a_{k+1} - a_k$  is same every time. Therefore,  $d = a$

The given numbers are in A.P.

Three more terms are

$$a_5 = 4a + a = 5a$$

$$a_6 = 5a + a = 6a$$

$$a_7 = 6a + a = 7a$$

**Q.18** The following series is APs? If its a form an A.P. find the common difference d and write three more terms.  $\sqrt{2}, \sqrt{8}, \sqrt{18}, \sqrt{32} \dots$

**Sol:**

$$\sqrt{2}, \sqrt{8}, \sqrt{18}, \sqrt{32} \dots$$

It can be observed that

$$a_2 - a_1 = \sqrt{8} - \sqrt{2} = 2\sqrt{2} - \sqrt{2} = \sqrt{2}$$

$$a_3 - a_2 = \sqrt{18} - \sqrt{8} = 3\sqrt{2} - 2\sqrt{2} = \sqrt{2}$$

$$a_4 - a_3 = \sqrt{32} - \sqrt{18} = 4\sqrt{2} - 3\sqrt{2} = \sqrt{2}$$

i.e.,  $a_{k+1} - a_k$  is same every time.

Therefore, the given numbers are in A.P. and  $d = \sqrt{2}$

Three more terms are

$$a_5 = \sqrt{32} + \sqrt{2} = 4\sqrt{2} + \sqrt{2} = 5\sqrt{2} = \sqrt{50}$$

$$a_6 = 5\sqrt{2} + \sqrt{2} = 6\sqrt{2} = \sqrt{72}$$

$$a_7 = 6\sqrt{2} + \sqrt{2} = 7\sqrt{2} = \sqrt{98}$$

**Q.19** The following series is APs? If its a form an A.P. find the common difference d and write three more terms.  $\sqrt{3}, \sqrt{6}, \sqrt{9}, \sqrt{12} \dots$

**Sol:**

$$\sqrt{3}, \sqrt{6}, \sqrt{9}, \sqrt{12} \dots$$

It can be observed that

$$a_2 - a_1 = \sqrt{6} - \sqrt{3} = \sqrt{3 \times 2} - \sqrt{3} = \sqrt{3}(\sqrt{2} - 1)$$

$$a_3 - a_2 = \sqrt{9} - \sqrt{6} = 3 - \sqrt{6} = \sqrt{3}(\sqrt{3} - \sqrt{2})$$

$$a_4 - a_3 = \sqrt{12} - \sqrt{9} = 2\sqrt{3} - \sqrt{3 \times 3} = \sqrt{3}(2 - \sqrt{3})$$

i.e.,  $a_{k+1} - a_k$  is not same every time.

Therefore, the given numbers are not in A.P.

**Q.20** The following series is APs? If its a form an A.P. find the common difference d and write three more terms.  $1^2, 3^2, 5^2, 7^2 \dots$

**Sol:**

$$1^2, 3^2, 5^2, 7^2 \dots \text{ or } 1, 9, 25, 49 \dots$$

It can be observed that

$$a_2 - a_1 = 9 - 1 = 8$$

$$a_3 - a_2 = 25 - 9 = 16$$

$$a_4 - a_3 = 49 - 25 = 24$$

i.e.,  $a_{k+1} - a_k$  is not the same every time.

Therefore, the given numbers are not in A.P.

**Q.21** The following series is APs? If its a form an A.P. find the common difference d and write three more terms.  $1^2, 5^2, 7^2, 73 \dots$

**Sol:**  $1^2, 5^2, 7^2, 73 \dots$  or  $1, 9, 25, 49 \dots$

It can be observed that

$$a_2 - a_1 = 25 - 1 = 24$$

$$a_3 - a_2 = 49 - 25 = 24$$

$$a_4 - a_3 = 73 - 49 = 24$$

i.e.,  $a_{k+1} - a_k$  is same every time.

Therefore, the given numbers are in A.P.

And,  $d = 24$

Three more terms are

$$a_5 = 73 + 24 = 97$$

$$a_6 = 97 + 24 = 121$$

$$a_7 = 121 + 24 = 145$$

**Q.22** Find two numbers whose sum is 27 and product is 182.

**Sol:** Let the first number be  $x$  and the second number is  $27 - x$ .

Therefore, their product =  $x(27 - x)$

It is given that the product of these numbers is 182.

$$\therefore x(27 - x) = 182$$

$$\Rightarrow x^2 - 27x + 182 = 0$$

$$\Rightarrow x^2 - 13x - 14x + 182 = 0$$

$$\Rightarrow x(x - 13) - 14(x - 13) = 0$$

$$\Rightarrow (x - 13)(x - 14) = 0$$

Either  $x - 13 = 0$  or  $x - 14 = 0$

i.e.,  $x = 13$  or  $x = 14$

If first number = 13, then

$$\text{Other number} = 27 - 13 = 14$$

If first number = 14, then

$$\text{Other number} = 27 - 14 = 13$$

Therefore, the numbers are 13 and 14.

**Q.23** Find two consecutive positive integers, sum of whose squares is 365.

**Sol:** Let the consecutive positive integers be  $x$  and  $x + 1$ .

$$\therefore x^2 + (x + 1)^2 = 365$$

$$\Rightarrow x^2 + x^2 + 1 + 2x - 365 = 0$$

$$\Rightarrow 2x^2 + 2x - 364 = 0$$

$$\Rightarrow x^2 + 13x + 14x - 182 = 0$$

$$\Rightarrow x(x + 13) + 14(x + 13) = 0$$

$$\Rightarrow (x + 13)(x + 14) = 0$$

Either  $x + 14 = 0$  or  $x - 13 = 0$ , i.e.,  $x = -14$  or  $x = 13$

Since the integers are positive,  $x$  can only be 13.

$$\therefore x + 1 = 13 + 1 = 14$$

Therefore, two consecutive positive integers will be 13 and 14.

**Q.24:** The altitude of a right triangle is 7 cm less than its base. If the hypotenuse is 13 cm, find the other two sides.

**Sol:** Let the base of the right triangle be  $x$  cm.

Its altitude =  $(x - 7)$  cm

$$\therefore x^2 + (x - 7)^2 = 13^2$$

$$\Rightarrow x^2 + x^2 + 49 - 14x = 169$$

$$\Rightarrow 2x^2 - 14x - 120 = 0$$

$$\Rightarrow x^2 - 7x - 60 = 0$$

$$\Rightarrow x^2 - 12x + 5x - 60 = 0$$

$$\Rightarrow x(x - 12) + 5(x - 12) = 0$$

$$\Rightarrow (x - 12)(x + 5) = 0$$

Either  $x - 12 = 0$  or  $x + 5 = 0$ , i.e.,  $x = 12$  or  $x = -5$

Since sides are positive,  $x$  can only be 12.

Therefore, the base of the given triangle is 12 cm and the altitude of this triangle will be  $(12 - 7)$  cm = 5 cm.

**Q.25** A cottage industry produces a certain number of pottery articles in a day. It was observed on a particular day that the cost of production of each article (in rupees) was 3 more than twice the number of articles produced on that day. If the total cost of production on that day was Rs 90, find the number of articles produced and the cost of each article.

**Sol:** Let the number of articles produced be  $x$ .

Therefore, cost of production of each article = Rs  $(2x + 3)$

It is given that the total production is Rs 90.

$$\therefore x(2x + 3) = 90$$

$$\Rightarrow 2x^2 + 3x - 90 = 0$$

$$\Rightarrow 2x^2 + 15x - 12x - 90 = 0$$

$$\Rightarrow x(2x + 15) - 6(2x + 15) = 0$$

$$\Rightarrow (2x + 15)(x - 6) = 0$$

Either  $2x + 15 = 0$  or  $x - 6 = 0$ , i.e.,  $x = -15/2$  or  $x = 6$ .

As the number of articles produced can only be a positive integer,  
therefore,  $x$  can only be 6.

Hence, number of articles produced = 6

Cost of each article =  $2 \times 6 + 3 = \text{Rs } 15$